

## 1118 Binary Stirling Numbers

The Stirling number of the second kind  $S(n, m)$  represents the number of ways to partition a set of  $n$  things into  $m$  nonempty subsets. For example, there are seven ways to split a four-element set into two parts:

$$\{1, 2, 3\} \cup \{4\}, \{1, 2, 4\} \cup \{3\}, \{1, 3, 4\} \cup \{2\}, \{2, 3, 4\} \cup \{1\}, \\ \{1, 2\} \cup \{3, 4\}, \{1, 3\} \cup \{2, 4\}, \{1, 4\} \cup \{2, 3\}.$$

We can compute  $S(n, m)$  using the recurrence,

$$S(n, m) = mS(n-1, m) + S(n-1, m-1), \text{ for integers } 1 < m < n.$$

but your task is slightly different: given integers  $n$  and  $m$ , compute the parity of  $S(n, m)$ , i.e.  $S(n, m) \bmod 2$ .

### Example

$$S(4, 2) \bmod 2 = 1.$$

Write a program that reads two positive integers  $n$  and  $m$ , computes  $S(n, m) \bmod 2$ , and writes the result.

### Input

The input begins with a single positive integer on a line by itself indicating the number of the cases following, each of them as described below. This line is followed by a blank line, and there is also a blank line between two consecutive inputs.

The input consists two integers  $n$  and  $m$  separated by a space, with  $1 \leq m \leq n \leq 1000000000$ .

### Output

For each test case, the output must follow the description below. The outputs of two consecutive cases will be separated by a blank line.

The output should be the integer  $S(n, m) \bmod 2$ .

### Sample Input

1

4 2

### Sample Output

1