

12805 Raiders of the Lost Sign

For many many years, mathematicians all over the world have focused their attention on one of the most fascinating phenomenon of Maths: the relationship between the perimeter and the diameter of a circumference (π).

Using one of Euler's formulas, we can calculate π as an addition of infinite terms,

$$\pi = 1 + 1/2 + 1/3 + 1/4 - 1/5 + 1/6 + 1/7 + 1/8 + 1/9 - 1/10 + 1/11 + 1/12 - 1/13 + \dots$$

Some of these terms add on, and others subtract, according to the following rules,

- You put '+' if the denominator of the fraction is 2.
- You put '+' if the denominator of the fraction is a prime like $4m - 1$, for some integer m .
- You put '-' if the denominator of the fraction is a prime like $4m + 1$, for some integer m .
- If the denominator is not a prime number, you put the sign resultant of multiplying the signs corresponding to each prime factor.

For example, the term with denominator 3 has '+' because 3 is prime and $3 = 4 \times 1 - 1$. The term with denominator 13 has '-' because 13 is prime and $13 = 4 \times 3 + 1$. The term with denominator 6 has '+' because $6 = 2 \times 3$, and 2 and 3 have respectively '+'. And the term with denominator 10 has '-' because $10 = 2 \times 5$, and 2 has '+' and 5 has '-'. So, 50 is $5 \times 5 \times 2$, i.e., $- \times - \times + = +$.

We want to obtain the sign of the i -th term of the former Euler's formula, i.e., the sign of the term with denominator i .

Input

The first line of the input contains an integer, t , indicating the number of test cases. For each test case, one line appears containing an integer i , $2 \leq i \leq 100000$.

Output

For each test case the output should contain a single line, indicating the sign of the term $1/i$ of Euler's formula (+ or -).

Sample Input

```
6
2
13
45
87
88
100000
```

Sample Output

+
-
-
-
+
-